

When Do Children and Teens Need Vaccinations?

Age	HepB Hepatitis B	DTaP/Tdap Diphtheria, tetanus, pertussis (whooping cough)	Hib <i>Haemophilus influenzae</i> type b	IPV Polio	PCV13 Pneumococcal conjugate	RV Rotavirus	MMR Measles, mumps, rubella	Varicella Chickenpox	HepA Hepatitis A	HPV Human papillomavirus	Men- ACWY	MenB	Influenza Flu
											Meningococcal		
at Birth (before hospital discharge)	✓												
2 months	✓	✓	✓	✓	✓	✓							
4 months	✓	✓	✓	✓	✓	✓							
6 months	✓	✓		✓	✓								✓
12 months							✓ (12 mos)	✓ (12 mos)					
15 months			✓ (15 mos)		✓ (15 mos)				✓✓				
18 months		✓ ² (18 mos)							(2 doses given 6 mos apart at age 18 & 24 mos)				
24 months													
5 years		✓		✓			✓	✓					
7–10 years													
11–12 years		✓ (Tdap)								✓✓✓	✓		
13–15 years													
16–18 years											✓	✓✓	

(One dose each fall or winter to all people ages 6 mos and older. Some children younger than age 9 years need 2 doses; ask your child's health-care provider if your child needs more than 1 dose.)

RED CHECK MARK AT 2, 4 AND 6 MOS. IS A SINGLE SHOT COMBINATION PEDIARIX.
MAGENTA CHECK MARK AT 5 YRS. IS A SINGLE SHOT COMBINATION KINRIX.
GREEN CHECK MARK AT 5 YRS. IS A SINGLE SHOT COMBINATION PROQUAD.