



## Acetaminophen (Tylenol)

Tylenol can be given every **4-6 hours** as needed for fever greater than or equal to 100.4 degrees. **Do not** give Acetaminophen to infants younger than 8 weeks old without consulting your pediatrician.

| Weight (Pounds) | Infant and Children's Liquid (160mg/5mL) | Chewable Tablets 80 mg | Junior Strength 160 mg |
|-----------------|--|------------------------|------------------------|
| 6-7             | 1.25 ml                                  |                        |                        |
| 8-9             | 1.75 ml                                  |                        |                        |
| 10-11           | 2 ml                                     |                        |                        |
| 12-13           | 2.5 ml (1/2 tsp)                         |                        |                        |
| 14-17           | 3 ml                                     |                        |                        |
| 18-23           | 3.75 ml                                  | 1 ½ tabs               |                        |
| 24-35           | 5 ml                                     | 2 tabs                 | 1 tab                  |
| 36-47           | 7.5 ml                                   | 3 tabs                 | 1 ½ tabs               |
| 48-59           | 10 ml                                    | 4 tabs                 | 2 tabs                 |
| > 60            | Follow instructions on box               |                        |                        |

## Ibuprofen (Motrin, Advil)

Ibuprofen can be given every **6-8 hours** as needed for fever greater than or equal to 100.4 degrees. **Do not** use in infants below 6 months of age or less than 16 lbs without consulting your pediatrician.

| Weight (Pounds) | Infant Drops (50mg/1.25mL) | Children's Liquid (100mg/5mL) | Chewable Tablets 50 mg | Chewable Tablets 100 mg |
|-----------------|----------------------------|-------------------------------|------------------------|-------------------------|
| 12-17           | 1.25 ml                    | 2.5 ml                        |                        |                         |
| 18-23           | 1.875 ml                   | 4 ml                          |                        |                         |
| 24-35           | 2.5 ml                     | 5 ml                          | 2 tabs                 |                         |
| 36-48           |                            | 7.5 ml                        | 3 tabs                 |                         |
| 48-59           |                            | 10 ml                         | 4 tabs                 | 2 tabs                  |
| >60             | Follow instructions on box |                               |                        |                         |

## Benadryl (Diphenhydramine)

| Weight (Pounds)              | 16-21  | 22-29 | 30-44    | 45-99  | >99         |
|------------------------------|--------|-------|----------|--------|-------------|
| Children's Liquid 12.5mg/5ml | 2.5 ml | 5 ml  | 7.5 ml   | 10 ml  |             |
| Chewable tablets 12.5 mg     | -      |       | 1 ½ tabs | 2 tabs | 4 tabs      |
| Capsules 25 mg               | -      | -     | -        | -      | 2 caps/tabs |