



## OUR VACCINATION SCHEDULE

Birth:	Hepatitis B
2 month visit:	Pediarix, Vaxneuvance, Pedvaxhib, Rotateq
3 month visit:	None
4 month visit:	Pediarix, Vaxneuvance, PedvaxHib
5 month visit:	None
6 month visit:	Pediarix, Vaxneuvance, Rotateq
12 month visit:	MMR, Varivax, Vaxneuvance
15 month visit:	DTaP, Pedvaxhib, Hepatitis A
2 year visit:	Hepatitis A
4 year visit:	Kinrix
5 year visit:	MMR, Varivax
11 year visit:	Tdap
12 year visit:	Menveo, Gardasil
16 year visit:	Menveo, Bexsero

\*Starting at 6 months patients are able to receive the annual influenza vaccine. The first year a patient receives the influenza vaccine they will receive 2 doses separated by at least 4 weeks. Every year thereafter they will need to receive only 1 dose.

\*If the first dose of Gardasil is given before 15yrs, patient will only need one additional dose at least 6 months later. If the first dose is given after 15yrs, patient will require a second dose at least 2 months after first and a third dose at least 4 months after second.

\*A second dose of Bexsero is needed 1 month after the first to complete the series.

Pediarix = DTaP (includes whooping cough), Hepatitis B, Polio

Vaxneuvance = Pneumococcus (bacteria)

PedvaxHib = Haemophilus influenza (bacteria)

Rotateq = Rotavirus

Varivax = Varicella (chickenpox)

Kinrix = DTaP, Polio

Menveo = Meningococcal Meningitis (types A,C,Y,W-135)

Bexsero = Meningococcal Meningitis (type B)

Gardasil = Human Papilloma Virus