

JOSE RIVAS DOMINGUEZ, M.D., LLC. OBSTETRICS • GYNECOLOGY • INFERTILITY

CIRCUMCISIONS

All circumcisions will be performed prior to you being discharged from the hospital, provided there are no medical problems and the clearance from the pediatrician is obtained.

If you are a self-pay patient or have Medicaid coverage you are required to make financial arrangements with our office prior to delivery. Circumcisions are not a covered benefit with Medicaid. The price is \$350.00.

Thank you for your attention in this matter.

Patient Name	Witness
	1
Signature	Date



JOSE RIVAS DOMINGUEZ, M.D., LLC. OBSTETRICS • GYNECOLOGY • INFERTILITY

Genetic Testing

AFP - Plus is maternal serum (blood) screen for birth defects. The AFP - Plus testing incorporates the features of AFP screening but detects three times as many cases of Down Syndrome as AFP testing alone.

AFP – Plus is a combination of three maternal serum (blood) testing; alphal fetoprotein (AFP), unconjugated estriol (uEST), and human chorionic gonadotrophin (HCG). The AFP – Plus screening procedure is based on studies showing that pregnancies affected with Down Syndrome have lower serum levels of AFP and uEST and higher levels of HCG than women with unaffected pregnancies.

AFP – Plus testing detects approximately 85% of open neural tube defects and approximately 60-70% of Down Syndrome pregnancies. AFP – Plus does not reliably screen for other chromosomal abnormalities.

AFP - Plus is a screening test. A normal result does NOT guarantee a normal baby. Should the AFP - Plus results be abnormal, additional diagnostic test or procedures may be recommended.

I have read the above and understand that benefits of AFP – Plus screening.

I request the following testing:

☐ AFP – Plus		i
☐ I decline any testing		
Patient Signature	Date	
·		
Witness	Date	



JOSE RIVAS DOMINGUEZ, M.D., F.A.C.O.G. OBSTETRICS • GYNECOLOGY • INFERTILITY

MaterniT 21 Plus- (Integrated Genetics Labcorp CONSENT FORM

MaterniT 21 plus prenatal test is a new type of test that analyzes DNA in a sample of your blood to predict the risk of Down syndrome (trisomy 21) and certain other genetic conditions.

MaterniT 21 plus prenatal test requires a single blood draw. It can be done as early as 10 weeks or later in pregnancy.

Other commonly used tests for Down syndrome are performed later in pregnancy and may require multiple office visits.

MaterniT 21 Plus Prenatal Test is a screening test that is more accurate than traditional Down syndrome blood tests. The MaterniT 21 Plus test is much less likely to give a false-positive result compared to traditional tests such as the first trimester screening test. That means there will be much less chance your doctor would recommend follow-up diagnostic testing, such as amniocentesis. 2

MaterniT 21 Plus also tests for two other genetic conditions, trisomy 18 (Edward syndrome) and trisomy 13 (Patau syndrome).

In addition, with MaternitT 21 Plus you have the option to evaluate fetal sex, X and Y sex chromosomes.

Non-invasive prenatal testing based on cell-free DNA analysis is not considered diagnostic. Once you have your Progenity Inc. test results, you can discuss your pregnancy care with your healthcare provider.

FINANCIAL RESPONSIBILITY

MaterniT 21 Plus is covered by many insurance plans. While the exact cost depends on your insurance coverage, Integrated Genetics Labcorp provides insured parents with a program that allows them to pay no more than a maximum out-of-pocket. If you do not have health insurance, or have financial need, there are additional MaterniT 21 Plus programs in the US that can assist with test costs.

Call 1-844-799-3243 for help with billing or payment questions.

Website: integratedgenetics.com/transparency

CPT: 81507 (NIPT) ,81599 (Fetal Sex)

Self-Pay Price: \$299.00 (Trisomy 13, 18, 21 & Fetal Sex) Test Code 451927



JOSE RIVAS DOMINGUEZ, M.D., F.A.C.O.G. OBSTETRICS • GYNECOLOGY • INFERTILITY

*Only patients that participation in the program qualify for the discounted price please call 1-844-799-3243 and simply register for the program.

You have 3 months to do survey and they survey will be emailed to you.

Inheritest Core Panel CPT 451964 \$299.00 max. Per study.

Inheritest Society Guided Panel CPT 451960 max. \$299.00 per study.

I hereby understand and authorize the office of my physician Jose Rivas Dominguez, MD to submit my blood collection for the MaterniT21plus test and understand that I must contact Integrated Genetic LabCorp to determine the financial cost of such test. I understand that in the event that my insurance carrier denies payment for the services I will be fully responsible for any fees associated with such test. Our office will not provide any medical necessity documentation to your insurance carrier for optional testing.

PATIENT CONSENT STATEMENT

I have read or have had read to me the above informed consent information about the MaternitT 21 Plus I have had the opportunity to ask questions of my health care provider regarding this test, including the reliability of test results, the risks, and the alternatives prior to giving my informed consent. I understand that my personal health information and my blood samples will be sent to Progenity Inc. I request and authorize Integrated Genetic LabCorp to test my sample(s) for the chromosome conditions listed above. I acknowledge that Integrated Genetics Lab Corp will send the results to my ordering healthcare provider. In the event of a high risk result, I acknowledge that Integrated Genetics LabCorp may contact my healthcare provider to obtain follow-up diagnostic information to ensure quality and accuracy in reporting. I acknowledge that I must sign the consent statement located on the test requisition form that will be sent with my sample(s) to Integrated Genetics LabCorp understand that I must also sign this consent form which will remain in my electronic medical chart.

Patient Signature:	Date:
<u> </u>	



JOSE RIVAS DOMINGUEZ, M.D., F.A.C.O.G.

OBSTETRICS • GYNECOLOGY • INFERTILITY

Printed Name:	DOB:
Witness/MD Signature:	



JOSE RIVAS DOMINGUEZ, M.D., F.A.C.O.G. OBSTETRICS • GYNECOLOGY • INFERTILITY

PROGENITY CONSENT FORM

Progenity Inc. Prenatal Test is a new type of test that analyzes DNA in a sample of your blood to predict the risk of Down syndrome (trisomy 21) and certain other genetic conditions.

Progenity Inc. Prenatal Test requires a single blood draw. It can be done as early as 10 weeks or later in pregnancy.

Other commonly used tests for Down syndrome are performed later in pregnancy and may require multiple office visits.

The Progenity Inc. Prenatal Test is a screening test that is more accurate than traditional Down syndrome blood tests. The Progenity Inc. test is much less likely to give a false-positive result compared to traditional tests such as the first trimester screening test. That means there will be much less chance your doctor would recommend follow-up diagnostic testing, such as amniocentesis.

Progenity Inc. also tests for two other genetic conditions, trisomy 18 (Edward syndrome) and trisomy 13 (Patau syndrome).

In addition, with Progenity Inc. you have the option to evaluate fetal sex, X and Y sex chromosomes.

Non-invasive prenatal testing based on cell-free DNA analysis is not considered diagnostic. Once you have your Progenity Inc. test results, you can discuss your pregnancy care with your healthcare provider.

FINANCIAL RESPONSIBILITY

Progenity Inc. is covered by many insurance plans. While the exact cost depends on your insurance coverage, Progenity Inc. provides insured parents with a program that allows them to pay no more than a maximum out-of-pocket. If you do not have health insurance, or have financial need, there are additional Progenity Inc. programs in the US that can assist with test costs.

Call 1-855-293-2639 for help with billing or payment questions.

CPT: 81507 (NIPT) ,81599 (Fetal Sex)

Self-Pay Price: \$549.00 Innatal (Trisotomy 13, 18, 21 & Fetal Sex)

\$349.00 Trio (CF, SMA and FragileX)

\$599.00 Standard (Screens 29 Disorders)



JOSE RIVAS DOMINGUEZ, M.D., F.A.C.O.G. OBSTETRICS • GYNECOLOGY • INFERTILITY

I hereby understand and authorize the office of my physician Jose Rivas Dominguez, MD to submit my blood collection for the Progenity Inc. test and understand that I must contact Progenity Inc. to determine the financial cost of such test. I understand that in the event that my insurance carrier denies payment for the services I will be fully responsible for any fees associated with such test. Our office will not provide any medical necessity documentation to your insurance carrier for optional testing.

PATIENT CONSENT STATEMENT

I have read or have had read to me the above informed consent information about the Progenity Inc. I have had the opportunity to ask questions of my health care provider regarding this test, including the reliability of test results, the risks, and the alternatives prior to giving my informed consent. I understand that my personal health information and my blood samples will be sent to Progenity Inc. I request and authorize Progenity Inc. to test my sample(s) for the chromosome conditions listed above. I acknowledge that Progenity Inc. will send the results to my ordering healthcare provider. In the event of a high risk result, I acknowledge that Progenity Inc. may contact my healthcare provider to obtain follow-up diagnostic information to ensure quality and accuracy in reporting. I acknowledge that I must sign the consent statement located on the test requisition form that will be sent with my sample(s) to Progenity Inc. understand that I must also sign this consent form which will remain in my electronic medical chart.

Patient Signature:	_Date:
Printed Name:	_DOB:
Witness/MD Signature:	



JOSE RIVAS DOMINGUEZ, M.D., F.A.C.O.G. OBSTETRICS • GYNECOLOGY • INFERTILITY

HARMONY CONSENT FORM

Harmony Prenatal Test is a new type of test that analyzes DNA in a sample of your blood to predict the risk of Down syndrome (trisomy 21) and certain other genetic conditions.

Harmony Prenatal Test requires a single blood draw. It can be done as early as 10 weeks or later in pregnancy.

Other commonly used tests for Down syndrome are performed later in pregnancy and may require multiple office visits.

The Harmony Prenatal Test is a screening test that is more accurate than traditional Down syndrome blood tests. The Harmony test is much less likely to give a false-positive result compared to traditional tests such as the first trimester screening test. That means there will be much less chance your doctor would recommend follow-up diagnostic testing, such as amniocentesis.²

Harmony also tests for two other genetic conditions, trisomy 18 (Edward syndrome) and trisomy 13 (Patau syndrome).

In addition, with Harmony you have the option to evaluate fetal sex, X and Y sex chromosomes.

Non-invasive prenatal testing based on cell-free DNA analysis is not considered diagnostic. Once you have your Harmony test results, you can discuss your pregnancy care with your healthcare provider.

FINANCIAL RESPONSIBILITY

Harmony is covered by many insurance plans. While the exact cost depends on your insurance coverage, Harmony provides insured parents with a program that allows them to pay no more than a maximum out-of-pocket. If you do not have health insurance, or have financial need, there are additional Harmony programs in the US that can assist with test costs.

Call 855-927-4672 for help with billing or payment questions.

CPT: 81507 (NIPT), 81599 (Fetal Sex)

Self- Pay Price: \$ 349.00

Should your insurance deny payment the fee per Harmony/ Ariosa will be \$199.00



JOSE RIVAS DOMINGUEZ, M.D., F.A.C.O.G. OBSTETRICS • GYNECOLOGY • INFERTILITY

I hereby understand and authorize the office of my physician Jose Rivas Dominguez, MD to submit my blood collection for the Harmony test and understand that I must contact Ariosa Lab to determine the financial cost of such test. I understand that in the event that my insurance carrier denies payment for the services I will be fully responsible for any fees associated with such test. Our office will not provide any medical necessity documentation to your insurance carrier for optional testing.

PATIENT CONSENT STATEMENT

I have read or have had read to me the above informed consent information about the Harmony. I have had the opportunity to ask questions of my health care provider regarding this test, including the reliability of test results, the risks, and the alternatives prior to giving my informed consent. I understand that my personal health information and my blood samples will be sent to Ariosa Diagnostics. I request and authorize Ariosa Diagnostics Lab to test my sample(s) for the chromosome conditions listed above. I acknowledge that Ariosa Diagnostics will send the results to my ordering healthcare provider. In the event of a high risk result, I acknowledge that Ariosa Diagnostic may contact my healthcare provider to obtain follow-up diagnostic information to ensure quality and accuracy in reporting. I acknowledge that I must sign the consent statement located on the test requisition form that will be sent with my sample(s) to Ariosa Diagnostics understand that I must also sign this consent form which will remain in my electronic medical chart.

Patient Signature:	Date:		
Printed Name:	DOB:		
Filmed Name.			
Witness/MD Signature:			



OBSTETRICS • GYNECOLOGY • INFERTILITY

CONSENT CARRIER TESTING FRAGILE X SYNDROME

The most common inherited cause of mental retardation is Fragile X syndrome. Symptoms involve developmental delay, mental retardation, autism and hyperactivity. Women who are carriers are at risk to have a child with mental retardation. Fragile X syndrome affects primarily boys and approximately 1 in 260 women. It can also occur in all ethnicities.

Inheritance

If the mother is a carrier, there is a 50% chance the child will have Fragile X syndrome.

Consent

If I am a carrier, prenatal testing is available to find out whether or not the baby has inherited the abnormal Fragile X gene.

Referral

We will arrange a consult with a Perinatologist for genetic counseling and additional testing if needed based on your results.

You should be certain you understand the following:

- The purpose of these tests is to determine whether I am a carrier of one of the common genetic abnormalities that cause Fragile X syndrome.
- The tests do not detect all carriers of the disease.
- The laboratory needs accurate information about my family history for the most accurate interpretation of the test results.
- The decision to have the carrier testing is completely mine.
- No other tests will be performed or reported on my sample unless authorized by my doctor. Any
 unused portion of my original sample will be destroyed within two months of receipt of the sample by
 the laboratory.
- The laboratory will disclose the results ONLY to my doctor, or to his/her agent, unless otherwise authorized by me or required by law.

I want Fragile X carrier testing	I do not want Fragile X carrier testing
Patient Signature	 ' Date
Witness	 Date



JOSE RIVAS DOMINGUEZ, M.D., LLC. OBSTETRICS • GYNECOLOGY • INFERTILITY

CONSENT CARRIER TESTING SPINAL MUSCULAR ATROPHY

Spinal Muscular Atrophy (SMA) is the most common inherited cause of early childhood death. It affects 1 in 35 to 1 in 117 people in the U.S. and varies by ethnicity. SMA destroys nerve cells that affect voluntary movement. Infants with SMA have problems breathing, swallowing, controlling head or neck movements, crawling and walking. The most common form affects infants in the first month of life and can cause death between 2-4 years of age. Less commonly the condition starts later and people can survive into adulthood. SMA does not affect the learning process and there is no cure or treatment.

Inheritance

If the test shows you are a carrier of SMA, the next step is for your partner to have carrier testing performed. Both parents must be carriers for the baby to be at risk for SMA. If your partner has a negative test result and no family history of SMA the chance of your baby having SMA is less than 1%. If both parents are carriers, there is 25% or a chance of 1 in 4 to have a child with SMA.

Referral

We will arrange a consult with a perinatologist for genetic counseling and additional testing if needed based on your results.

You should be certain you understand the following:

- The purpose of these tests is to determine whether I am a carrier of one of the common genetic abnormalities that cause SMA.
- The tests do not detect all carriers of the disease.
- The laboratory needs accurate information about my family history for the most accurate interpretation of the test results.
- The decision to have the carrier testing is completely mine.
- No other tests will be performed or reported on my sample unless authorized by my doctor. Any
 unused portion of my original sample will be destroyed within two months of receipt of the sample by
 the laboratory.
- The laboratory will disclose the results ONLY to my doctor, or to his/her agent, unless otherwise authorized by me or required by law.

I want SMA carrier testing	I do not want SI	MA carrier testing
		1
Patient Signature		Date
Witness	·	Date:



JOSE RIVAS DOMINGUEZ, M.D., LLC: OBSTETRICS • GYNECOLOGY • INFERTILITY

PROCEDURE: CESAREAN SECTION

Your doctor has determined that it will be necessary to deliver your baby by Cesarean section or "C-section." This is a major operation involving a surgical cut through the abdominal or "belly" wall and through the wall of the "womb." Complications from this type of delivery are uncommon, but they do occur. Therefore, your doctor makes no guarantee as to the possible result obtained from method of delivery.

As with a surgical procedure, bleeding and infection are possible complications of C-section. These complications can result in prolonged illness, permanent deformity, poorly healing wounds, the need for blood transfusions, and the need for further major surgery. On rare occasions, it is necessary for the doctor to remove the womb during C-section to stop bleeding. Damage to the urinary system and the intestines are also rare complications of C-sections. This type of damage can result in malfunction of the intestines and the female organs. A hernia or "rupture" developing at the site of the surgical cut in the belly is another uncommon complication of this operation. Because there will be a scar in the womb, the chance of rupturing during future pregnancy is higher.

Some of the complications of C-section may require further surgery; some may cause permanent deformity; some can result in sterility or the permanent inability to become pregnant; rarely some can even be fatal. However, many of the above complications are also associated with vaginal deliveries. Therefore, in those patients in whom C-section is indicated the procedure may provide the patient; with the best chance of success and the lowest risk of complications.

ADDITIONAL RISKS AND	ALTERNATIVES:		1
		ssary):	i .
I CERTIFY: I have read or involved in this procedu questions have been answ	re; I have had the o	contents of this form; I understand the opportunity to ask any questions which	e risks and alternatives th I had and all of my
DATE:	TIME:	SIGNED:(Signed by the person legally a	uthorized to consent for patient)
WITNESS:			
WITH 110001			ĺ



OBSTETRICS • GYNECOLOGY • INFERTILITY

HIV TEST IN PREGNANCY . CONSENT FORM

Information is provided in accordance to Florida Law

HIV/AIDS is an important health concern for pregnant women because she can pass the HIV virus to her baby during pregnancy or childbirth or through breastfeeding. HIV testing is recommended as a routine test for all pregnant women. It is much better for a woman to know her HIV status as early in pregnancy as possible so she can make important decisions about health care and breast feeding. Tests are available to detect antibodies for HIV that are safe and can be done along with other prenatal blood tests.

A positive test does not necessarily mean that you have AIDS or that you will become ill with AIDS. A positive test does mean that can infect others with the virus and that you must take precautions to prevent spreading the infection. If your test is positive, you will again knowledge and understanding of an important medical condition and be able to inform your sexual partner (s) and health care provider (s).

There are medications that may help a woman who is pregnant and has HIV to reduce the chance of passing HIV to her baby. If a pregnant woman is HIV-positive and does not get treatment, her baby has about a 25% chance of getting HIV from her. But if an HIV-positive pregnant woman receives appropriate medication as late as during the delivery of her child, she can reduce the risk of transmission by at least 50%.

A negative test result may mean that you have not been infected with HIV-1. If you have been engaging in behaviors that put you at risk, you may want to be retested in approximately six months. A negative test may also mean that your body has not had time to develop antibodies to HIV-1 and that you have an early infection.

Because treatment is so effective in preventing babies from getting HIV, Florida law and regulations require that every pregnant woman be counseled about HIV and the benefits of testing and be offered and HIV test along with the standard blood test for syphilis and hepatitis B surface antigen (HBsAg). Testing must be offered at the time of the first examination relating to the current pregnancy and again at 28 to 32 weeks gestation.

Although HIV testing is routinely performed as part of the antenatal testing protocol, you have the right to refuse the test. The decision to have testing for syphilis, hepatitis B, or HIV is voluntary and you may withdraw your consent at any time.

Your physician will answer any questions you may have about HTV testing. If you are pregnant and you test positive for HTV, your physician can provide the care you need and information about services and options available to you. Your physician can tell you the risk of passing HTV infection to your baby, about medications given during pregnancy that can significantly reduce the risk of passing HTV to your baby, and the medical care available for babies who may be infected with HTV.



OBSTETRICS • GYNECOLOGY • INFERTILITY

CONSENT TO HIV-1 ANTIBODY TESTING IN PREGNANCY

The purpose of the test, its potential uses, and the limitations and the meaning of the rules have been explained to me. I understand that if the results indicate that if the results indicate that my blood contains antibody to HIV, it means that I

may ha	have been infected with the HIV virus, which is believed to caus	e AIDs (Acquired Immune Deficiency Syndrome).		
AT FIRS	FIRST PRENATAL VISIT			
	I authorize my healthcare providers to collect one or more blood specimen form me at the time of my first prenatal visit in order to detect whether or not I have antibodies in my blood to HIV- 1 (human immunodeficiency virus). This is the virus which has been associated with AIDS (Acquired Immune Deficiency Syndrome). I understand that my physician will report test results to me in person and not by telephone or by mail. At that time, I will have the opportunity to receive counseling about the meaning of the test results, the possible need for retesting, and other matters. Information regarding measures for the prevention of, exposure to, and transmission of HIV has been made available to me.			
Lundo	sent to <u>Release</u> derstand that the test results will be confidential and will not be nitted or required by law, I also consent to the release of the test	e disclosed to any person without my consent unless results to		
	REFUSAL OF HIV-1 ANTIBODY TESTING With the information presented above having been explain understand, all of my questions having been answered and give my consent for HIV testing.	ned to me completely and clearly in the language I with full knowledge of the consequences, I refuse to		
Patient	ient Signature	Date/Time		
Witnes	tness	Name of Patient (Please Print)		
<u>IN TH</u>	THIRD TRIMESTER			
	Authorization for Repeat HIV Testing in Third Trimester of I authorize my health care provider to repeat the testing for pregnancy. This consent for repeat testing is limited to the my health care provider will discuss testing with me before the test results.	e course of my current pregnancy. I understand that		
	I Decline Repeat HIV Testing in Third Trimester of Pregna With the information presented above having been expla understand, all of my questions having been answered ar repeat testing for sexually transmitted diseases and HIV la	and with full knowledge of the consequences, I decline		
—— Patie	atient Signature	Date/Time		
— Witn		Name of Patient (Please Print)		



OBSTETRICS • GYNECOLOGY • INFERTILITY

CONSENT TO CARRIER TESTING CYSTIC FIBROSIS

Cystic Fibrosis (CF) is an inherited disease that affects more than 25,000 American children and young adults. Symptoms of CF vary but include lung congestion, pneumonia, diarrhea, and poor growth. Most people with CF have severe medical problems and some die at a young age. Others experience few symptoms and may be unaware of having CF, there is no cure for CF at this time and it does not affect the learning process in children. Many people with CF have died at a very young age in the past. As a result of scientific advantages these days many with the disease are living into their 20's and 30's.

WHAT ARE THE CHANCES OF MY BABY HAVING CYSTIC FIBROSIS?

You can have a child with CF even if there is no family history. Carrier frequency in the U.S. is 1 in 30 on average and varies by ethnicity. CF testing can help determine if you are a carrier and at risk to have two altered genes. Most people have two normal copies of the CF gene.

You should be certain you understand the following:

- The purpose of these tests is to determine whether I am a carrier of one of the common genetic abnormalities that cause CF.
- The tests do not detect all carriers of the disease.
- The laboratory needs accurate information about my family history for the most accurate interpretation of the test results.
- The decision to have the carrier testing is completely mine.
- No other tests will be performed or reported on my sample unless authorized by my doctor. Any
 unused portion of my original sample will be destroyed within two months of receipt of the
 sample by the laboratory.
- The laboratory will disclose the results ONLY to my doctor, or to his/her agent, unless otherwise authorized by me or required by law.

I want CF carrier testing		I do not want CF carrier	testing	
Patient Signature	_			Date
			1	
Witness			• 1	Date

Cord Blood Stem Cell Preservation Frequently Asked Questions

- What is cord blood? Cord blood is the blood remaining in your child's umbilical cord following birth. It is a rich, non-controversial source of stem cells that must be collected at the time of birth.
- What are stem ceils? Stem cells are the building blocks of our blood and immune systems. They are found throughout the body including bone marrow, cord blood and peripheral blood. They are particularly powerful because they have the ability to treat, repair and/or replace damaged cells in the body.
- 3. Why do families choose to bank their newborn's stem cells? Today, cord blood stem cells have been used in the treatment of approximately 80 diseases. Banking your baby's stem cells guarantees the cells will be available to your family should you need to use them. Cord blood is also being used in emerging treatments, for diseases like Type 1 Diabetes² and Cerebral Palsy, which require a child's own cord blood. Stem cells from a related source are the preferred option for all treatments, and transplants using cord blood from a family member are more likely to be successful than transplants using cord blood from a non-relative (i.e., a public source).
- 4. How is cord blood collected? The collection process is safe, easy and painless for both mother and baby and does not interfere with the delivery. After the baby is born, but before the placenta is delivered, a medical professional will clean a 4 to 8 inch area of the umbilical cord with antiseptic solution and insert a needle connected to a blood bag into the umbilical vein. The blood flows into the bag, by gravity until the umbilical vein is emptied. The blood bag is clamped, sealed, labeled and shipped by courier to a processing lab. The collection itself typically takes about 2 to 4 minutes.
- 5. Who can use my newborn's stem cells? Your newborn's stem cells have the potential to be used by the child, and, if there's an adequate match, by siblings and sometimes parents. An adequate match using related cord blood is defined as a 3 of 6 HLA Match. When two people share the same HLAs, they are said to be a 'match' which means their tissues are immunologically compatible. Your newborn's cord blood will always be a 100% match for him or herself and there is up to a 75% probability of a match for a sibling.⁵

Options for Cord Blood & Cord Tissue Banking:

200	ecting a child is exciting for both new and e sions to make for your family's future, inclu on that's right for you. (Check one below):	experienced parents. You ding choosing a cord blo	I have many important ood and cord tissue banking
	☐ Family Banking Family banking allows families to save to their family's use. This service is availab to collect, process, freeze and store your for your family's future medical use. Cor	le through private cord bl r child's stem cell-rich um	ood banks that charge a fee
	☐ Donation Cord blood donation is a way for you to p the blood of your child's umbilical cord in donate the cord blood to a public facility blood. Find out if your hospital accepts o donated.)	na donation facility for the vour family does not ret	e public good. Once you
	☐ Discard Discard umbilical cord blood and tissue a	s waste. The cells cannot	t be retrieved for future use.
have health	e read the information above and discussence provider. I have indicated my choice	d my cord blood and tissi above.	ue banking options with my
		•	
Patier	nt Signature		Date
	For Medical Profe	essional's Use Only	
Pa	atient Name:	DOB:	
М	edical Professional:	Cord Blood Bank:	
			

There is no guarantee that banked cord blood or tissue will be a match for a family member or will be an appropriate or effective treatment.

Autologous cord blood and cord tissue stem cells will not guarantee suitable treatment for all inherited genetic diseases.