



Martin A. Garcia, MD ▪ **Erika D. Glas, DO**
Tiffany A. Wells, MD ▪ **Kendra K.H. Gillespie, MD**
Diplomates, American College of Obstetrics and Gynecology
Jan Ely, APRN ▪ **Sheri Ray, APRN** ▪ **Molly Turner, APRN**
Claudette Mallory, APRN ▪ **Michelle Buoye, APRN**

Baptist Medical Pavilion ▪ 836 Prudential Drive ▪ Suite 1103 ▪ Jacksonville, FL 32207
Satellite ▪ 9889 Gate Parkway ▪ Suite 205 ▪ Jacksonville, FL 32246
Phone (904) 398-7654 ▪ Fax (904) 398-0118

DENTAL GUIDELINES IN PREGNANCY

Our OB patients are encouraged to continue their routine preventative dental care. It is recommended to defer care that is considered strictly cosmetic until after delivery.

- 1) Local Anesthetic agents are satisfactory, preferably without Epinephrine. Epinephrine would be acceptable if bleeding is a significant concern.
- 2) It is recommended to avoid inhalation agents or heavy sedation during procedures unless anesthesiology monitoring is utilized. We would consider procedures under general anesthesia safe when administered by an anesthesiologist in the appropriate setting.
- 3) X-rays can be performed as indicated with the use of abdominal shielding.
- 4) Left lateral decubitus positioning may help avoid syncopal episodes.
- 5) Approved Antibiotics: Erythromycins, Penicillin's, Cephalosporin's, Augmentin & Zithromax

Contraindicated Antibiotics: Tetracycline's and Quinolones

- 6) Approved Pain Medicine: Acetaminophen

Contraindicated Pain Medicine: Ibuprofen products and other NSAIDs
After 13 weeks gestation, these can be used for less than 5 days duration

These are our general guidelines for medications and antibiotics; however we will be glad to assist in reviewing any proposed medication use from pregnancy perspective.

Please call our office with any questions.

Thank you,
The River City OBGYN Team